

## ENHANCEMENT OF MACROPHAGE TUMOURICIDAL ACTIVITY BY THE ALKALOID DERIVATIVE UKRAIN. *IN VITRO* AND *IN VIVO* STUDIES.

SOTOMAYOR E.M.<sup>1</sup>, RAO K.<sup>1</sup>, LOPEZ D.M.<sup>1</sup>, LIEPINS A.<sup>2\*</sup>

- 1) Department of Microbiology and Immunology, University of Miami School of Medicine, Miami, Florida, USA.
- 2) Division of Basic Medical Sciences, Faculty of Medicine, Memorial University of Newfoundland, St. John's, Newfoundland, A1B 3V6 Canada.

**Summary:** *Ukrain is a semisynthetic drug with immunomodulatory properties, derived from Chelidonium majus L. alkaloids and thiophosphoric acid. The effect of this compound on the growth of Balb/c syngenic mammary adenocarcinoma was assessed. Intravenous, but not subcutaneous or intraperitoneal, administration of this drug was found to be effective in delaying tumour growth in an actual therapeutic protocol initiated five days after tumour implantation. No untoward side-effects were observed using these in vivo treatment modalities. The role of macrophages in the observed retardation of tumour development was investigated using peritoneal exudate macrophages (PEM) in cytotoxicity assays. In previous studies, the authors have found that PEM of mammary tumour bearing mice lose their capacity to kill a variety of tumour target cells including the in vitro cultured homologous tumour cells (DA-3). Pretreatment of PEM from normal mice with 2.5  $\mu$ M Ukrain for 24 h followed by stimulation with either IFN- $\gamma$  or with LPS + IFN- $\gamma$  enhanced their cytotoxic activity. Treatment of PEM from tumour bearing mice with 2.5  $\mu$ M Ukrain and LPS results in a reversal of their defective cytotoxic response against the DA-3 target cells. Furthermore, Ukrain alone, in the absence of a secondary signal, induced the activation of tumouricidal function of PEM from tumour bearing but not from normal mice. These data indicate that Ukrain's in vivo effects against the development of mammary tumours may be due, at least in part, to its ability to restore macrophage cytolytic function.*

### Introduction

The use of various biological response modifiers (BRMs) which enhance the host immune response, is an active area of investigation due to their possible clinical applications (1-3). The full therapeutic potential of the various existing BRMs has not been fully realized because of their undesirable side effects when used at concentrations required for maximum biological activity (4-6). Thus, the identification and characterization of new compounds that

might increase the immunological activity without deleterious side effects, would enhance the prospects of immunotherapy as a practical and effective cancer treatment modality.

A new and promising area of research has evolved from biochemical phytotherapy, where plant alkaloids, as well as their semisynthetic derivatives (phytopharmaceuticals), have been found to be effective immunomodulators in animal models (7-9) and in cancer patients (10, 11). For example, the indolizidine alkaloid swainsonine has been reported to inhibit experimental lung metastasis of B16-F10 melanoma cells when administered systemically

\* Author to whom correspondence should be addressed.

to C57BL/6 mice in a prophylactic setting. The inhibition of pulmonary metastasis was found to be mediated by the host's NK cells (7, 8). Moreover, swainsonine induced an increase in murine splenocytes and macrophage cytotoxicity and proliferation of bone marrow cells as well as HLA Class I expression on human breast carcinoma cells (9, 12).

Similarly, alkaloids from the plant *Chelidonium majus* L. conjugated to thiophosphoric acid yield a triaziridide compound denoted as Ukrain which, in preliminary clinical studies, has been found to have immunomodulatory activity (10, 11). The major immune parameters affected by this alkaloid derivative consist of an increase in the total T-lymphocyte number and a normalization of the T-helper/T-suppressor cell ratios, without affecting serum immunoglobulin levels, complement components and acute phase proteins (10).

The present studies were undertaken to investigate the possible mechanism by which Ukrain mediates its therapeutic effects (10, 11). For this purpose the authors chose the D1-DMBA-3 mammary tumour syngenic to Balb/c mice which in previous studies has been found to cause a profound impairment of various parameters of the immune system (13–20). *In vivo* studies utilizing this tumour system showed that intravenous administration of Ukrain is effective in reducing the rate of growth of established tumours. It is well known that macrophages (PEM) are important non-specific effector cells with antitumour activity (21, 22). The present *in vitro* results demonstrate that Ukrain is able to reverse the defective capacity of PEM from tumour bearing (TB) mice to lyse tumour targets upon LPS stimulation. Furthermore, the restored lytic activity is TNF- $\alpha$  independent, thus indicating that Ukrain activates an alternate lytic mechanism in macrophages from tumour bearing mice.

#### Materials and methods

**Mice and tumours.** BALB/c mice are maintained by brother-sister matings in the laboratory of the

Department of Microbiology and Immunology, Miami. The tumour D1 DMBA-3 is a transplantable mammary adenocarcinoma derived from a non viral, non carcinogen-induced preneoplastic alveolar nodule in a BALB/c mouse after treatment with 7,12-dimethylbenzanthracene (23). D1-DMBA-3 is non metastatic and immunogenic to the host of origin. The tumour is routinely transplanted in BALB/c by s.c. injection of  $1 \times 10^6$  tumour cells. The tumour becomes apparent five days after implantation; by day 30 necrotic areas are evident and the mice begin to die.

**Biological response modifier.** Ukrain is a semi-synthetic compound consisting of *Chelidonium majus* L. alkaloids conjugated to thiophosphoric acid. This compound is positively charged with m.w. of 1,470 and melting point of 222–223°C. The manufactured product, which complies with the good practice manufacture guidelines for pharmaceutical products, is available at 1.0 mg of Ukrain per 1.0 ml of H<sub>2</sub>O (J.W. Nowicky Pharmaceuticals, Margaretenstrasse 7, 1040 Vienna, Austria). In *in vitro* studies the authors used the pertinent tissue culture medium as drug diluent, and PBS in *in vivo* studies.

**In vivo treatment with Ukrain.** Ukrain treatment was initiated five days after subcutaneous tumour implantation. Three routes of administration were employed, i.e., intravenous, intraperitoneal and subcutaneous. All three experimental groups, of at least five mice each, received 4.0  $\mu$ g Ukrain in 0.1 ml of PBS. This dosage was chosen based on preliminary experiments. The treatment protocol was five consecutive injections at 24 h intervals, followed by 72 h without the drug. This regimen was repeated three times, i.e., three weeks. The mean tumour diameters were determined on days 7, 9, 12, 14, 16, 19, 21 and 23 after tumour implantation. The data, i.e., mean tumour diameters, were statistically analysed by a two-way repeated measures analysis of variance for differences between routes of drug (Ukrain) delivery at eight successive time points. The four averages (three experimental and one control)

at each time point were compared by Tukey's w-procedure at  $p < 0.05$ . The sample size of five animals for each of four groups re-measured eight times gave sufficient degrees of freedom ( $= 17$  for between groups residual error) for a precise estimation of the between group residual mean square ( $SD^2$ ) in ANOVA.

**Reagents.** The culture medium used in these assays was RPMI 1640 supplemented with 100 units of penicillin and 100  $\mu\text{g/ml}$  of streptomycin,  $5 \times 10^5$  M 2-ME, 2 mM L-glutamine, 1% non-essential amino acids, 1% essential amino acids, 1% sodium pyruvate (all from GIBCO Laboratories, Grand Island, NY), and 10% endotoxin-free fetal bovine serum (Hyclone Laboratories, Logan UT). LPS (*E. coli* 055:B5) from DIFCO Laboratories, Detroit, MI and murine  $\text{INF-}\gamma$  from Genzyme Corporation, Boston, MA were used as indicated.

**Target cells.** DA-3 cell line was derived from the *in vivo* D1-DMBA-3 mammary tumour syngenic to BALB/c mice. The DA-3 cell line produces tumours in BALB/c mice with the same growth kinetics and expresses the same tumour associated antigens on its surface as the parent tumour. The DA-3 cell line was maintained *in vitro* as cell suspension in RPMI 1640 FBS supplemented media.

**Macrophage cultures.** Normal mice and three- to four-week tumour bearers were injected i.p. with 1 ml of thioglycolate (DIFCO Laboratories, Detroit, MI). On day 4, the PEM were obtained by peritoneal lavage with 10 ml ice-cold RPMI 1640/mouse. Peritoneal cells were washed twice and re-suspended in supplemented RPMI 1640 media. The adherent population was obtained following the plastic-adherence technique described by Penhale (24). This procedure provides a population consisting of greater than 95% macrophages as determined by staining with Diff-Quick differential stain (Baxter, McGraw Park, IL) and nonspecific esterase staining (Sigma). Viability of cell was routinely  $>95\%$  by trypan blue exclusion. PEM were seeded into 96-well, flat bottom microtitre plates (Costar, Cambridge, MA) at cell densities required for the particular assay.

**Macrophage cytotoxicity assay.** Cytotoxicity was determined by the release of  $^{51}\text{Cr}$  from labelled tumour target cells. Briefly, purified PEM from normal mice and tumour bearers were first incubated in either medium alone or with different concentrations of LPS and/or  $\text{INF-}\gamma$ . After 24 h, macrophage monolayers were washed twice with warm RPMI 1640, and  $5 \times 10^3$  labelled target cells were seeded/well. Labelled tumour cells were prepared by incubating them with 0.1  $\text{mCi}^{51}\text{Cr}$  (New England Nuclear, Boston, MA) for 1 h, after which they were washed three times and resuspended in supplemented RPMI 1640 medium. The cultures were further incubated for 16 h and then 0.1 ml of culture medium was removed from each well, and the amount of radioactivity determined in an automatic gamma counter (Packard Auto-Gamma 500, Santa Clara, CA.). Percent specific lysis was calculated by the formula:


$$\% \text{ specific lysis} = \frac{\text{experimental cpm} - \text{spontaneous cpm}}{\text{maximum cpm} - \text{spontaneous cpm}} \times 100$$

where experimental cpm = counts released in wells containing macrophages and tumour cells; spontaneous cpm = counts released by target cells cultured alone; maximum cpm = the amount of  $^{51}\text{Cr}$  released by totally lysed target cells.

## Results and discussion


Ukrain has previously been found to have therapeutic potential (10, 11); however, its mode of action in tumour bearing subjects is largely unknown. To understand the underlying mechanism by which this compound exerts its anti-tumour effect, *in vivo* and *in vitro* studies were performed using a well defined mouse mammary adenocarcinoma model system (23).

*In vivo* studies were performed to investigate whether Ukrain would have an effect on the development of the D1-DMBA-3 mammary tumour. Three different routes of drug delivery were tested in order



**Fig.1** Effects of Ukrain (4.0 µg/mouse) administered subcutaneously (▨) or intraperitoneally (▩), on the *in vivo* growth of DA-3 mammary adenocarcinoma tumours in Balb/c mice. These routes of drug administration had no inhibitory effect on the tumour growth rate when compared to controls (■). At day 21 mice began to die due to tumour burden, hence the apparent differences in tumour size are not statistically significant.

to evaluate possible differences in therapeutic efficacy. As shown in Fig.1, neither subcutaneous nor intraperitoneal injections of Ukrain (4 µg/mouse) had a tumour growth inhibitory activity. In sharp contrast were the results obtained when Ukrain (4 µg/mouse) was given intravenously to tumour bearing mice. As shown in Fig. 2, significant ( $p < 0.05$ ) cessation of tumour growth was first detected on the fifth day after the start of periodic chemotherapy, i.e., fourteen days after tumour inoculation. Tumour growth continued to progress in the control mice and in treated mice; however, the rate of growth was significantly diminished in the latter. It should be emphasized that in these studies an actual therapeutic protocol was used, i.e., the administration of the drug was not initiated until four days after tumour implantation. Furthermore, neither the experimental nor control mice receiving Ukrain showed any deleterious drug related side effects. In preliminary and ongoing studies using higher doses of Ukrain (8 µg/mouse) and following the same protocol described above, a striking therapeutic effect has been observed in that



**Fig.2** Effects of Ukrain (4.0 µg/mouse) administered intravenously (□) on the *in vivo* growth of established DA-3 mammary adenocarcinoma tumours in Balb/c mice. This route of drug administration produced significant ( $p < 0.05$ ) inhibition of tumour growth at day 14 and thereafter as compared to control mice (■) in the experiment received five consecutive injections at intervals, followed by 72 h without drug. This regimen was repeated three times. None of the treated mice, regardless of route of administration, showed any adverse effects.

only one of five mice treated with this dosage developed tumours at day 15, while all five control mice had tumours and began to show signs of cachexia.

Previous *in vitro* studies with Ukrain showed that this compound is capable of activating the lytic function of lymphocytes obtained from the spleen of alloimmunized mice in a dose dependent manner (25). At 1.2 µM this drug was found to increase several fold the cytolytic activity of the spleen lymphocytes obtained from C57BL/6 (H-2<sup>b</sup>) mice immunized with mastocytoma P815 (H-2<sup>d</sup>) cells in a 3.5 h <sup>51</sup>Cr release assay at an E/T ratio of 5:1. The enhanced lytic activity was dose dependent and remained specific for the cells used as immunogen (25). Hence, the present studies were designed to investigate whether Ukrain has a similar effect on macrophage population in normal mice and in tumour bearing animals. Initial studies were performed using PEM from normal mice, which were treated

Table I Effects of Ukrain on the capacity of PEM from normal mice to lyse tumour cells in response to LPS and/or INF- $\gamma$

Pretreatment (24 h) Ukrain	% Specific Cytotoxicity			
	None	Activation (24 h)		
		LPS <sup>1</sup>	INF- $\gamma$ <sup>2</sup>	LPS + INF- $\gamma$ <sup>3</sup>
0	0	10.5 $\pm$ 1.8	2.9 $\pm$ 2.0	14.0 $\pm$ 4.1
25.0 $\mu$ M	0	0	0	0
5.0 $\mu$ M	5.0 $\pm$ 0.7	0	0	3.1 $\pm$ 0.9
2.5 $\mu$ M	2.4 $\pm$ 1.4	12.0 $\pm$ 2.6	7.0 $\pm$ 2.6	20.1 $\pm$ 1.9

E/T ratio: 60:1

<sup>1</sup>LPS: 10  $\mu$ g/ml

<sup>2</sup>INF- $\gamma$ : 5.0 U/ml

<sup>3</sup>LPS 10  $\mu$ g/ml + INF 5.0 U/ml

with various concentrations of Ukrain for 24 h, followed by stimulation with LPS and/or INF- $\gamma$  for an additional 24 h. As shown in Table I, Ukrain alone, when used at low concentrations, was able to trigger a nominal macrophage cytotoxicity against syngenic DA-3 mammary tumour cells. However, Ukrain at 25  $\mu$ M and 5  $\mu$ M did not prime PEM from normal animals to become more responsive to subsequent stimulation with either LPS, INF- $\gamma$ , or combination thereof. In sharp contrast were the results obtained with Ukrain used at 2.5  $\mu$ M. This dose was able to prime macrophages to display enhanced cytotoxic activities when they were further stimulated with either 5 U/ml of INF- $\gamma$  (2.9  $\pm$  2.0 vs 7.0  $\pm$  2.6% killing) or a combination of LPS + INF- $\gamma$  (14.0  $\pm$  4.1 vs 20.1  $\pm$  1.9% killing). Thus, 2.5  $\mu$ M of the drug seemed to be the optimal concentration needed to achieve an enhancing effect on macrophage tumouricidal activity. Based on these results, this concentration of drug was used in subsequent experiments.

The progressive growth of the D1-DMBA-3 mammary adenocarcinoma in BALB/c mice results in a profound impairment of the capacity of PEM obtained from these animals to kill a variety of tumour target cells upon *in vitro* stimulation with LPS. Significantly, the syngenic DA-3 mammary tumour cells

Table II Effects of Ukrain on the cytolytic activity of PEM from normal and tumour bearing mice

Treatment	Normal PEM % lysis	PEM from tumour-bearing mice % lysis
None	0	0
Ukrain 2.5 $\mu$ M <sup>1</sup>	4.8 $\pm$ 1.7	13.0 $\pm$ 3.1
LPS 10 $\mu$ g/ml <sup>2</sup>	29.0 $\pm$ 2.0	12.0 $\pm$ 1.7
LPS 10 $\mu$ g/ml + Ukrain	25.0 $\pm$ 3.7	24.0 $\pm$ 3.0
INF- $\gamma$	0	0
INF- $\gamma$ + Ukrain <sup>3</sup>	4.5 $\pm$ 0.9	11.0 $\pm$ 0.8
LPS + INF- $\gamma$	31.0 $\pm$ 3.0	37.0 $\pm$ 1.2
LPS + INF- $\gamma$ + Ukrain	39.0 $\pm$ 3.0	39.0 $\pm$ 2.0

are not killed by macrophages from mammary tumour bearing animals (19). Thus, subsequent experiments were designed to investigate the conditions that might overcome the inability of these PEM from tumour bearing animals to kill tumour cells *in vitro*. As illustrated in Table II, incubation of PEM from normal and tumour bearers with 2.5  $\mu$ M of Ukrain alone for 24 h resulted in tumouricidal activation of PEM from tumour bearing mice (13%), whereas PEM from normal mice did not display such significant activation (4.8%). Moreover, Ukrain added together to LPS did not alter the cytolytic capacity of normal mice PEM in response to LPS activation. In contrast, this alkaloid was able to overcome the defective response of PEM from tumour bearers to LPS stimulation (from 12% to 24%). In recent experiments the authors have found that this increase in lytic activity is associated with a significant decrease in TNF- $\alpha$  production, i.e., from 50 down to 5.0 U/ml (data not shown). These results suggest that immunostimulation by Ukrain, resulting in reversal of the depressed macrophage cytotoxicity in tumour bearers, involves a TNF- $\alpha$  independent mechanism of action.

Previous studies have shown that INF- $\gamma$  used in combination with other agents can activate

**Table III** Effects of Ukrain on the cytolytic activity of PEM from normal and *in vivo* LPS-treated mice

<i>In vitro</i> treatment (24 h)	PEM from normal % lysis	PEM from LPS-treated mice % lysis
None	0	0
Ukrain 2.5 $\mu$ M	4.8 $\pm$ 1.7	8.2 $\pm$ 1.1
LPS 10 $\mu$ g/ml	29.0 $\pm$ 1.0	0
LPS + Ukrain < 2.5 $\mu$ M	25.0 $\pm$ 3.7	23.5 $\pm$ 2.3
INF- $\gamma$ 5 U/ml	0	0
INF- $\gamma$ + Ukrain 2.5 $\mu$ M	4.5 $\pm$ 0.9	16.2 $\pm$ 2.3

macrophage tumouricidal activity (26). As shown in Table II, incubation of PEM from normal and tumour bearers with 5 U/ml of INF- $\gamma$  alone, did not result in triggering of macrophage cytotoxicity against DA-3 targets. Moreover, no enhancement (over that induced by Ukrain alone) was noted when this cytokine was used in combination with Ukrain, either in PEM from normal or in PEM from tumour bearing mice. In addition, Ukrain added to the combination of LPS and INF- $\gamma$  did not result in a significant further enhancement of macrophage tumouricidal activities over those obtained with LPS + INF- $\gamma$  alone. These results suggest that Ukrain is exerting its immune enhancing effect at the level of LPS-inducible mechanism of macrophage cytotoxicity.

To evaluate further whether this drug can also reverse the depressed macrophage cytotoxicity found in other experimental conditions, the authors chose the *in vivo* LPS desensitization model for study. LPS desensitization is a well known phenomenon that occurs when normal mice and/or macrophages are treated either *in vivo* or *in vitro* with a small dose of LPS followed by a second challenge with the same agent at higher doses (27). This treatment leads to inactivation of macrophage lytic function and cytokine production. As shown in Table III, PEM from *in vivo* LPS-treated mice (10  $\mu$ g/mouse 24 h before PEM harvest) could not be activated to kill tumour cells in response to a second *in vitro* challenge with LPS, as compared to PEM from untreated mice

which only received the activating agent *in vitro* (0% vs 29.0% killing). However, when Ukrain was added together with LPS *in vitro*, this treatment resulted in a reversal of the suppressed cytotoxicity of PEM from LPS-treated mice, thus overcoming the *in vivo* LPS-induced desensitization. Moreover, this drug when used alone, was able to induce higher levels of macrophage cytotoxicity in PEM from LPS-treated mice than in PEM from control untreated mice when it was used alone (8.2 vs 4.8% killing) or when used in combination with INF- $\gamma$  (16.2% vs 4.5% killing).

The data available from previous studies (10, 11, 25) taken in conjunction with the results presented herein indicate that Ukrain functions as an up-regulator of the immune system and as an effective antitumour drug when administered *i.v.* to tumour bearing mice. Not only are the various effector cells functionally enhanced by Ukrain, but in preliminary studies the authors have found a decrease in the *in vitro* TNF- $\alpha$  production by macrophages treated with this drug. Since the authors have observed that the tumour bearing mice die after development of cachexia, it is tempting to speculate that an additional possible benefit of Ukrain treatment could be the amelioration of the cachectic state.

#### Acknowledgements

These studies were supported by grants R37 CA22583 and RO1 CA54226 from the National Institutes of Health, USA. The excellent technical assistance of Mantley Dorsey, Jr. and the excellent secretarial services of Ms. Doris Williams are greatly appreciated. Dr. David Bryant provided the learned advice necessary for the statistical analysis and interpretation of the *in vivo* data.

#### References

- (1) Terry W.D., Rosenberg S.A. (eds.) Immunotherapy of Human Cancer: Elsevier North-Holland, New York 1982.
- (2) Herberman R.B. *Biological response modifiers for the*

- therapy of cancer. In: Tsubura E.I., Urushizaki, A., Sonne T., Kameko Y. (eds.) "Rationale of Biological Modifiers in Cancer Treatment." Excerpta Medica, Princeton, 1985, pp. 240-255.
- (3) Rosenberg S.A., Lotze M.T. *Cancer immunotherapy using Interleukin-2 and Interleukin-2 activated lymphocytes.* Annu. Rev. Immunol., 4, 681-709, 1986.
- (4) Lotze M.T., Frana L.W., Sharrow S.O., Robb R.J., Rosenberg S.A. *In vivo administration of purified human Interleukin 2. I. Half-life and immunologic effects of the Jurkat cell line-derived interleukin 2.* J. Immunol., 134, 157-166, 1985.
- (5) Lotze M.T. et al. *In vivo administration of purified human Interleukin 2. II. Half-life, immunologic effects, and expansion of peritoneal lymphoid cells in vivo with recombinant IL-2.* J. Immunol., 135, 2865-2875, 1985.
- (6) Urba W.J., Longo D.L.  *$\alpha$ -Interferon in the treatment of nodular lymphomas.* Semin. Oncol., 13 (Suppl. 5), 40-47, 1986.
- (7) Dennis J.W. *Effects of swainsonine and polyinosinic: polycytidylic acid on murine tumour cell growth and metastasis.* Cancer Res., 46, 5131-5136, 1986.
- (8) Humphries M.J., Matsumoto K., White S.L., Molyneux R.J., Olden K. *Augmentation of murine natural killer cell activity by swainsonine, a new antimetastatic immunomodulator.* Cancer Res., 48, 1410-1415, 1988.
- (9) Mohla S. et al. *Inhibition of growth of subcutaneous xenografts and metastasis of human breast carcinoma by swainsonine: modulation of tumor cell HLA class I antigens and host immune effector mechanisms.* Anticancer Res., 10, 1515-1522, 1990.
- (10) Nowicky J.W., Staniszewski A., Zbroja-Sontag W., Slezak B., Nowicky M., Hiesmayr W. *Evaluation of thiophosphoric-acid alkaloid derivatives from Chelidonium majus L. ("Ukrain") as an immunostimulant in patients with various carcinomas.* Drugs Exptl. Clin. Res., XVII, 139-143, 1991.
- (11) Nowicky J.W., Liepins A., Zbroja-Sontag W., Staniszewski A., Danilos J. *Evaluation of clinical studies of Ukrain in cancer patients.* Proc VII Mediterranean Congr. Chemotherapy., Barcelona, May, 1990.
- (12) Newlon S.A., White S.L., Humphries M.J., Olden K. *Swainsonine inhibition of spontaneous metastasis.* J. Natl. Cancer Inst., 81, 1024-1028, 1989.
- (13) Lopez D.M., Sigel M.M., Charyulu V. *Kinetics of responses to tumor antigens and mouse mammary tumor virus in BALB/cCrgt mice.* J. Natl. Cancer Inst., 66, 191-196, 1981.
- (14) Paul R.D., Ghaffar D., Sigel M.M., Charyulu V., Lopez D.M. *Splenic alterations during mammary tumorigenesis: diverse effects on different immune parameters.* Anticancer Res., 1, 63-69, 1989.
- (15) Bussow S.C., Paul R.D., Lopez D.M. *Influence of mammary tumor progression on phenotype and function of spleen and in situ lymphocytes in mice.* J. Natl. Cancer Inst., 73, 249-255, 1984.
- (16) Lopez D.M. et al. *Cytotoxic effector mechanisms involved in the immunity against mammary tumors.* In: Rich, M.A., Hager J.C., Lopez, D.M. (eds.) "Breast Cancer: Scientific and Clinical Progress". Kluwer Academic Publ., Boston, 1988, p. 185.
- (17) Fu Y.-X., Paul R., Wang Y., Lopez D.M. *Thymic involution and thymocyte phenotypic alterations induced by murine mammary adenocarcinomas.* J. Immunol., 143, 4300-4307, 1989.
- (18) Watson G.A., Fu Y.-X., Lopez D.M. *Splenic macrophages from tumor bearing mice co-expressing Mac-1 and Mac-2 antigens exert immunoregulatory functions via two distinct mechanisms.* J. Leukocyte Biol., 49, 126-138, 1991.
- (19) Sotomayor E.M., et al. *Role of tumor derived cytokines on the immune system of mice bearing a mammary adenocarcinoma. II. Downregulation of macrophage-mediated cytotoxicity by tumor-derived granulocyte-macrophage colony-stimulating factor.* J. Immunol., 147, 2816-2823, 1991.
- (20) Lopez D.M., Lopez-Cepero M., Watson G.A., Ganju A., Sotomayor E.M., Fu Y.-X. *Modulation of the immune system by mammary tumor-derived factors.* Cancer Invest., 9, 643-653, 1991.
- (21) Hibbs J.B., Lambert Jr L.H., Remington J.S. *Possible role of macrophage mediated nonspecific cytotoxicity in tumour resistance.* Nature, 235, 48-50, 1972.
- (22) Fidler I.J., Barnes Z., Fogler W.E., Kirsch R., Bugelski P., Poste G. *Involvement of macrophages in the eradication of established metastases following intravenous injection of liposomes containing macrophage activators.* Cancer Res., 42, 496-501, 1982.
- (23) Medina D., DeOme K.B. *Response of hyperplastic alveolar nodule outgrowth line D1 to mammary tumor virus, nodulo-inducing virus, and prolonged hormonal situation acting singly and in combination.* J. Natl. Cancer Inst., 42, 303-310, 1969.
- (24) Pennline K.J. *Adherence to plastic or glass surfaces.* In: Hercowitz, H.B., Holder, H.T., Bellanti, J.A., Ghaffar, A. (eds.) "Manual of Macrophage Methodology: Collection, Characterization and Function." Marcel Dekker, New York, 1981, p. 63.
- (25) Liepins A. *Enhancement of cell mediated lysis of tumor cells by Chelidonium majus L. alkaloids (Ukrain).* J. Cancer Res. Clin. Oncol., 116 (Suppl. 1), 436, 1990.
- (26) Pace J.L., Russell S.E., Torres B.A., Johnson H.M., Gray P.W. *Recombinant mouse  $\gamma$  interferon induces the priming step in macrophage activation for tumor cell killing.* J. Immunol., 130, 2011-2013, 1983.
- (27) Wallach D., Holtmann H., Engelmann H., Nophar Y. *Sensitization and desensitization to lethal effects of tumor necrosis factor and IL-1.* J. Immunol., 140, 2994-2999, 1988.